

# **On the Late Permian Age ( $258.3 \pm 2.5$ Ma) and Tectonic Significance of the Cataracte Pegmatitic Leucogranite (Valcea County, Romania)**

Mihai N. Ducea<sup>1,2,\*</sup>, Irina Tene<sup>2</sup>, Constantin Balica<sup>3</sup>, Ioan Balintoni<sup>3</sup>, and Horst Peter Hann<sup>4</sup>

<sup>1</sup> University of Arizona, Department of Geosciences, Tucson, AZ 85721, USA

<sup>2</sup> Faculty of Geology and Geophysics, University of Bucharest, 010041, Bucharest, Romania

<sup>3</sup> Department of Geology, Babes-Bolyai University of Cluj, 400084, Romania

<sup>4</sup> Institut für Geowissenschaften, Universität Tübingen, 72076 Tübingen, Germany

\*-corresponding author: [ducea@arizona.edu](mailto:ducea@arizona.edu)

## **Abstract**

We present new zircon U-Pb age data on a leucogranitic laccolith with pegmatitic textures from Cataracte (Valcea County, Romania). The Cataracte body is an S-type granite which was not deformed by regional metamorphism and intrudes into the Sebes Lotru Terrane (SLT) of the Dacia Super-terrane. SLT was metamorphosed to amphibolite facies during continental collision marking the Variscan orogeny (325-350 Ma) and is cross-cut by numerous pegmatitic dikes and sills of syn to late kinematic origin previously dated by zircon U-Pb at 317-325 Ma. In contrast, zircons from the Cataracte laccolith reveal a much younger, concordant,  $258.3 \pm 2.5$  Ma age. This late Permian age is coincident with the age of tectonic juxtaposition of the SLT against lower grade units along the Sibisel Shear Zone located some 40 km to the north and east in modern coordinates. We conclude that the intrusive rock at Cataracte is a rare magmatic body related to this little acknowledged cryptic collisional event that marks regionally the geologically important closure of Paleo-Tethys.

**Keywords:** Cataracte, Sebes-Lotru Terrane, leucogranite, Permian, Paleo-Tethys.

**Resume:**

Nous présentons de nouvelles données géochronologiques (méthode U-Pb sur zircons) de la mise en place d'une intrusion laccolitique de composition leucogranitique dans la région de Cataracte (Comté de Valcea en Roumanie). Le massif de Cataracte est un granite de type S intrusif au sein de la terrane de Sebes Lotru (SLT) et qui n'est pas affecté par la déformation et le métamorphisme régional. Appartenant à la Dacia Super-terrane, la SLT a quant à elle, été métamorphisée dans les conditions du faciès amphibolite au cours de la collision continentale associée à l'orogénèse varisque (325-350 Ma) et est recoupée par de nombreux dykes et sills pegmatitiques syn- à tardi-tectoniques (datés par U-Pb sur zircons à 317-325 Ma). En revanche, les zircons de l'intrusion laccolitique de Cataracte révèlent un âge concordant plus jeune de  $258.3 \pm 2.5$  Ma. Cet âge fini-Permien coïncide avec la phase d'empilement tectonique de la SLT sur les unités de bas grade, le long de la zone de cisaillement de Sibisel située 40 km au NE de la zone étudiée. Par conséquent, nous interprétons le massif intrusif de Cataracte comme un vestige rare de cet événement collisionnel resté cryptique dans la région, marquant la fermeture de la Paléo-Téthys.

**Mots-clés :** Cataracte, Sebes-Lotru, Terrane, leucogranite, Permien, Paléo-Téthys

## **Introduction**

This paper reports some new zircon U-Pb ages on a leucogranite laccolith-like body (hereafter referred to as the Cataracte body) with pegmatitic textures from the Sebes Lotru Terrane (SLT, Balintoni et al., 2014). Previously only a whole rock Pb-Pb isochron age of 333 Ma was reported from this general area in an abstract format (Ledru et al., 1997); the meaning and accuracy of this age or the location could not be evaluated since they were not formally published. Other syn to late kinematic dikes and sills of various pegmatites from the SLT (Hann, 1987 for a detailed early review) were later indeed dated to be of that age (317-325 Ma) by zircon U-Pb (Balintoni et al., 2010a, our own unpublished data). However, this particular igneous body has not been dated, even though it stands out as one of the largest leucogranitic (and pegmatitic) igneous body within the SLT. We show that its age is somewhat surprisingly late Permian and coincident with the development of an important ductile shear zone nearby, one that possibly marks the suture of the Paleo-Tethys regionally. Consequently, we speculate this newly determined age has regional tectonic implications which will be discussed below.

## Geologic background

The basement of Romanian Carpathians (Balintoni, 1997) is made of a various peri-Gondwanan Neo Proterozoic to early Paleozoic arc and back-arc terranes that were accreted to each other and to larger continental masses during the Paleozoic (Balintoni et al., 2014; Ducea et al., 2016). Most likely these terranes formed at the consuming margin of the Rheic ocean (von Raumer et al., 2008, 2013). During accretion they were metamorphosed and in places, were subject to more magmatism. They were further fragmented and shuffled during the Alpine orogeny (e.g. van Hinsbergen et al., 2020; Schmid et al., 2020 and references therein). Unlike in parts of the Alps, no high-grade Alpine metamorphism is superimposed on these Carpathian basement units; Alpine metamorphism is limited to local prehnite-pumpellyite facies in the South Carpathians (Ciulavu et al., 2008). For more in depth reviews of regional geology, the reader is referred to recent and comprehensive regional review papers by Matenco (2017), Iancu and Seghedi (2017), van Hinsbergen et al., (2020), Schmid et al., (2008, 2020) and references therein.

In detail, the geologic and especially the petro-tectonic evolution of Carpathian basement terranes continue to be poorly known although much progress has been made recently on deciphering ages of protoliths (Balintoni et al., 2009, 2010a, 2014) and metamorphism (Dragusanu and Tanaka, 1999, Medaris et al., 2003; Ducea et al., 2016). Still, correlations between various units and timing of their assembly are far from being resolved. The SLT (Figure 1) as the largest single coherent basement unit of the Romanian Carpathians, is located in the South Carpathians and occupies the bulk of the Sebes and Lotru Mountains as well as nearby ranges (Balintoni et al., 2014). It is part of the Getic nappe or thrust sheet (Iancu et al., 2005) which structurally lies on top of a Jurassic oceanic thrust sheet, the Severin (not pictured in Figure 1 a). The structure was developed in the earliest Cretaceous, although precise ages of thrusting have yet to be determined. Together with the units structurally above it, the SLT forms the Median Dacides of Sandulescu (1984), also known as Dacia Super-terrane (sometimes referred to as Dacia Mega-Unit). The SLT does have a latest Paleozoic unmetamorphosed cover in a few places and is also locally covered by some younger sedimentary rocks of the Alpine cycle (Iancu et al., 2005).

The SLT comprises a latest Precambrian (Ediacaran, 560-580 Ma) thin continental crustal basement exposed in the southern parts of the SLT, and named the *Lotru Metamorphic Unit* (Balintoni et al., 2010a). This thin arc basement unit was invaded by an arc formed at

about 440-480 Ma, with a high flux peak in the Mid-Ordovician at 466 Ma (Stoica et al., 2016, Balintoni et al., 2010a, Ducea et al., 2018) (Figure 1 b). This unit, which is dominated by mafic to intermediate volcanic arc successions (now metamorphosed) is known as the *Cumpana Metamorphic Unit* (Balintoni et al., 2010a). There are inherited zircon indications that this Ordovician-Silurian arc was formed off-shore the main Gondwanan mass but close to the present Arabian-Nubian shield. Consequently, the terrane is characterized by few zircons of ages older than 1 Ga. The latest magmatic manifestations of this peri-Gondwanan arc were in the earliest Silurian (440 Ma, the Cozia gneiss) and together, these two units were metamorphosed during the Carboniferous (Medaris et al., 2003).

Regional geology of the SLT in the central south Carpathians is more complex than the representation of the Lotru and Cumpana Metamorphic units depict in our text and Figure 1b. Other regional papers and geologic maps present a more complex division of units (e.g, Sabau, 1994, Stelea, 1994, Hann, 1995), but the simplicity of the Balintoni et al. (2010a) unit presentation- a latest Precambrian thin basement unit intruded by a massive transitional arc unit of Ordovician age – captures the bulk of SLT basement geology. One additional important regional unit is the Negovanu Mare formation (not pictured on our simplified map, but seen in the map of Stelea, 1999), a metasedimentary (mica schists, quartzites) unit, which may be lying unconformably over these arc terranes, hence it was probably deposited later. Nevertheless, the Negovanu Mare formation, as well as the rest of the Cumpana and Lotru units were metamorphosed together during the Variscan collisional orogeny.

In the SLT, regional metamorphism primarily of amphibolite facies is documented to be Carboniferous, as best determined by garnet 317-355 Ma Sm-Nd isochron ages (Medaris et al., 2003). Numerous meter- to decameter dikes and sills of S-type pegmatitic leucogranites or simply “pegmatites” (Hann, 1987) intrude and mostly cross-cut the metamorphic fabric. Their age has been determined to be around 317-325 Ma in a few places in the Sebes, Cibin and Lotru Mts (Balintoni et al., 2010a, Ducea, in progress, 2021) indicating that this age range seals the ductile deformation. While not properly interpreted as such in the literature, these leucogranites are syn to late collisional products marking the latest stages of compression that generated regional amphibolite facies metamorphism here. This age is widely recognized regionally to represent a Variscan event. It marks the timing of a Himalayan-style collision (Stern, 1994, von Raumer et al., 2013) that was followed in many places throughout Europe by extensional collapse in the earliest Permian (Menard and Molnar, 1988).

The Cataracte body (Manecan, 1985, Hann, 1987) is a poorly exposed pegmatitic leucogranite intruded into the Lotru Metamorphic Unit of the SLT in the southern part of the

Lotru Mts., some 10 km NW of the town of Voineasa (Valcea County), along the Lotru river (Savu and Schuster, 1975). Its best exposures are found in a region where the Lotru river cuts a short but deep gorge into this granitoid. The name of the body (meaning “waterfalls” in Romanian) comes from the waterfalls formed by the Lotru river in that area. The body has a laccolith like shape that is about 3x2 km in horizontal dimensions and about 0.8 km in vertical dimension. The overall shape of the body is determined by outcrop mapping (Savu and Schuster, 1975) but also from many km of mining adits and boreholes that were dug into the body for geologic prospecting and white mica mining work during the 1970s and 1980s in communist Romania.

At outcrop levels, the body contains numerous large enclaves of the metamorphic rocks nearby and is not as uniform as it may appear at larger scale; its appearance at small scale is more of a network of hundreds of sills intruded into the metamorphic basement which at the geologic map scale (Savu and Schuster, 1975) takes the shape of a laccolith. Detailed studies indicate some zoning at the scale of the intrusion, with an outer domain richer in mafic minerals (Hann, 1987). It is pegmatitic in most places, with crystal as large as many cm. Petrographically, it is a low temperature, S-type leucogranite dominated by quartz, k-feldspar commonly with myrmekitic textures, two micas, and almandinic garnet in places. Zircon, tourmaline and apatite are common accessory minerals; however, the intrusion is free of rare accessory minerals found commonly in pegmatites worldwide. The S-type character of this leucogranite is evident from whole-rock major, trace elements as well as high radiogenic isotopes (with measured  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios as high as 0.9) (Ducea et al., 2021, unpublished). In places, the Cataracte body appears concordant (even at large scale, hence its laccolithic appearance) but it is clearly not internally deformed. The great majority of outcrops display igneous textures that suggest that it clearly postdates the Variscan regional metamorphism.

### **Sample and methods**

Sample V15 (N45.465911/E23.871381) was collected from the core of the Cataracte body in an outcrop along the Lotru river. It is a fresh, coarse grained pegmatitic granitoid with major quartz, k-feldspar, plagioclase, muscovite and biotite visible in hand specimen and apatite, tourmaline and zircon found as accessory minerals and seen only in thin section. The studied sample is an S-type granite, which is significantly more radiogenic than most of the SLT (Ducea, unpublished data), due to the high Rb/Sr ratio, in turn constrained by the abundant micas and alkali-feldspars.

The zircons from the sample were extracted using the traditional method of crushing, sieving, heavy liquid separation, magnetic separation and hand-picking. The ~40 zircon grains were then mounted on epoxy with SL, R33 and FC zircon grains used as primary and secondary standards (see detailed method and analytical procedure in Pullen et al., 2018). Growth features and morphology of each zircon grains were identified using cathodoluminescence (CL) images using the Gatan ChromaCL2 scanning electron microscope in the Arizona LaserChron SEM Lab. Spot locations for analysis were then picked using the CL images to locate core and rim boundaries in complex zircons. In situ isotopic analysis of the zircons was conducted using the Thermo Element2 single-collector inductively coupled plasma mass spectrometry (ICPMS) coupled to a Photon Machines Analyte G2 Excimer Lasers at the Arizona LaserChron Center (ALC) (Gehrels et al., 2006, 2008; Gehrels and Pecha, 2014). The U-Pb age for each analysis was then calculated using the ALC in-house software E2agecalc, and for concordant subsets, a weighted average age was calculated using Isoplot 4.15 (Ludwig, 2008).

## Results

We measured 30 individual crystals from sample V15 (Table 1). Zircons from Cataracte are typically high U with values as high as 12% U, but commonly in the range of 2000-3000 ppm U. These zircons also have high U/Th ratios, typically in excess of 100. Only one measured crystal is inherited and had an age of 540 Ma, which is typical for the Lotru Metamorphic Unit of the SLT. Most crystals are inclusions-free, euhedral or subhedral and about 100-200 microns in their long dimension, suggesting a magmatic origin (despite the large U/Th).

Several of the analyzed grains are metamict, as seen in the secondary electron (SE) or backscattered (BSE) images taken on the SEM. Most of these crystals are not zoned in BSE images or luminescent in CL (Figure 2). A few non metamict crystals do show some complex crystallization zoning (Figure 2), however their age is the same as those of the metamict crystals and do not display any discernable age trends from core to rim.

Excepting the inherited age, the remainder of ages were evaluated on concordance. Those grains that are discordant were not further used in age calculation, whereas the 23 grains that passed the concordance test were pooled into a  $258.3 \pm 2.5$  Ma age with a mean of  $258 \pm 1.7$  Ma at a confidence level of 95% with a MSWD of 2.1. This is the first late Permian age of a granitoid in the SLT and in the South Carpathians in general. Similar age (250-270 Ma)

pegmatites are however described from the western Carpathians and Pannonian regions (Slovakia and Hungary) (Uher and Broska, 1996, Poler et al., 2002). Similar age granites are also found in the foreland east of the modern Carpathians in North Dobrogea (Balintoni et al., 2010b), but they represent a distinct geologic block not relatable to the Dacia Superterrane in particular.

## **Interpretations**

High U/Th ratios are rather common in South Carpathian pegmatites (Balintoni et al., 2010a) and probably owe this to sequestration of Th in the source in the form of monazite. However, unlike the smaller pegmatitic dikes and sills previously measured from the SLT, which were all dated at 317-325 Ma, the Cataracte body is much younger. Another significant difference is that the smaller dikes and sills have a large proportion of inherited grains among their zircon population (>50%), whereas in the case of Cataracte body, the great majority of analyzed zircons reveal their crystallization age. Only one out of 30 grains is inherited. Given the U-rich nature of the measured grains, it is not surprising that many of the crystals are metamict. However, given that most ages are concordant, we interpret that the pooled age of  $258.3 \pm 2.5$  Ma is geologically significant and represents the crystallization age of this body.

Is there a tectonic event to which this intrusion is to be connected? Recently, Ducea et al. (2016) have shown that the Sibisel Shear zone located some 40 km to the north and east (Figure 1b) was probably active as a major ductile structure in the latest Permian. This important steep compressional ductile structure (Pana and Erdmer, 1994, Hann, 1995) juxtaposed the SLT against the Rausorul Cisnadioarei Series, a greenschist facies meta andesite dominated unit that, unlike the SLT, was metamorphosed during the Ordovician (Ducea et al., 2016). The ages of mylonitization in the shear zone (near the village of Rasinari) are rather poorly constrained by Rb-Sr isochron ages and span the late Permian and possibly earliest Triassic (270-249 Ma) (Ducea et al., 2016). We speculate that the soft collision or transpressional event interpreted to have generated the Sibisel Shear Zone (Ducea et al., 2016) is responsible for the melting event that produced the Cataracte laccolith, since it is the only regional tectonic event of that age and is a process that commonly produces S-type granites by *self ignition* of high U metasedimentary rocks in thickened crust.

The larger significance of the Sibisel Shear Zone remains somewhat unresolved but it clearly juxtaposes two peri-Gondwanan terranes of similar igneous origin but very different

metamorphic histories prior to their assembly in the Permian. This is a likely candidate to represent a - or the suture of the Paleo-Tethys in the Romanian Carpathians.

*Acknowledgments.* M.N.D. acknowledges support from the Romanian Executive Agency for Higher Education, Research, Development and Innovation Funding project PN-III-P4-ID-PCCF-2016-0014. Ducea dedicates this paper to his undergraduate mentor, Prof. Emil Constantinescu, who as a principal investigator of a project set out to understand the Cataracte pegmatite in the early 1990s, engaged M.N.D. in his first research project as an undergraduate on rocks from there. Derek Hoffman is thanked for assistance with field and laboratory work, and Nicky Giesler for help with the SEM. Data are available as a Supplementary excel sheet and also available on the open science data platform at <https://osf.io/2grp6/>. This contribution has the following document identifier available through osf.io: DOI 10.17605/OSF.IO/2GRP6

### **Figure captions**

**Figure 1.** A. Major elements of the basement units in the South Carpathians (modified after Balintoni et al., 2010a). Terranes: SL- Sebes Lotru, L- Leaota, Fg-Fagaras, P-Pades, C-Caras, D- Danubian. Other elements: I-E Iuti Eibental ophiolite, M- various pre-Alpine un-metamorphosed magmatic rocks. Areas in white are younger rocks (Mesozoic and younger). This map contains no structural elements for simplicity. B. A more detailed map of the central part of the South Carpathians showing the location of the Cataracte body and the main geographic locations and geologic elements discussed in text. The outline of the Lotru and Cumpana metamorphic units is from Balintoni et al. (2010a). Only some minor later faults are shown in this figure, but no major tectonic elements of the Alpine fold and thrust belt, which is of no significance in this paper.

**Figure 2.** SEM images of metamict (left panel) and non-metamict (right panel) zircons from the Cataracte pegmatite. In each panel (a) represents a secondary electron (SE) photograph, (b) a backscattered (BS) image, and (c) a cathodoluminescence image. The SE and BS scales are identical.



**Figure 3.** (a). Concordia diagram for sample V15 with data point ellipses plotted at 2 sigma. (b). Best age for sample V15 based on  $^{238}\text{U}/^{206}\text{Pb}$  ages pooled among concordant grains. Drawn with a modified version of Isoplot 3.7 (Ludwig, 2008).

**Table1.** Zircon U-Pb data for sample V15, Cataracte body (provided as separate excel sheet, but also available at osf.io as described under the Acknowledgment section).

## References

Balintoni, I., 1997, Geotectonics of metamorphic terranes from Romania (in Romanian). Ed. Carpatica, 176 pp Cluj Napoca.

Balintoni, I., Balica, C., Ducea, M., Chen, F., Hann, H.P., Sabliovschi, V., 2009, Late Cambrian-Early Ordovician Gondwanan terranes in the Romanian Carpathians: a zircon U/Pb provenance study, *Gondwana Research* 16, 119–133, doi: 10.1016/j.gr.2009.01.007.

Balintoni, I., Balica, C., Ducea, M., Hann, H., Sabliovschi, V., 2010a, The anatomy of a Gondwanan terrane: The Neoproterozoic-Ordovician basement of the pre-Alpine Sebeş-Lotru composite terrane (South Carpathians, Romania), *Gondwana Research*, 17, 561-572, doi: 10.1016/j.gr.2009.08.003.

Balintoni, I., Balica, C., Seghedi, A., and Ducea, M. N., 2010b, Avalonian and Cadomian terranes in North Dobrogea, Romania: *Precambrian Research*, 182, 217-229, doi: 10.1016/j.precamres.2010.08.010

Balintoni, I., Balica, C., Ducea, M., Hann, H., 2014, Peri-Gondwanan terranes in the Romanian Carpathians: A review of their spatial distribution, origin, provenance and evolution, *Geoscience Frontiers*, 5, 395–411, doi: 10.1016/j.gsf.2013.09.002.

Ciulavu, M., Ferreira Mählmann, R., Schmid, S.M., Hofmann, H., Seghedi, A., Frey, M., 2008, Metamorphic evolution of a very low - to low-grade metamorphic core complex (Danubian window) in the South Carpathians. *Geol. Soc. London, Spec. Publ.* 298, 281—315.

Ducea, M.N., Negulescu, E., Profeta, L., Sabau, G., Jianu, D., Petrescu, L. and Hoffmann, D., 2016, Evolution of the Sibişel Shear Zone (South Carpathians): A study of its type locality near Răşinari (Romania) and tectonic implications. *Tectonics* 35, 9, 2131-2157, doi: 10.1002/2016TC004193.

Ducea, M. N., Giosan, L., Carter, A., Balica, C., Stoica, A. M., Roban, R. D., Balintoni, I., Filip, F., and Petrescu, L., 2018, U-Pb Detrital Zircon Geochronology of the Lower Danube and Its Tributaries: Implications for the Geology of the Carpathians: *Geochemistry Geophysics Geosystems*, 19, 9, 3208-3223, doi: 10.1029/2018GC007659.

Drăguşanu, C., Tanaka, T., 1999, 1.57-Ga magmatism in the South Carpathians: implications for the pre-Alpine basement and evolution of the mantle under the European continent. *Journal of Geology*, 107, 237-248.

Gehrels, G. and Pecha, M., 2014, Detrital zircon U-Pb geochronology and Hf isotope geochemistry of Paleozoic and Triassic passive margin strata of western North America: *Geosphere*, 10, 1, 49-65.

Gehrels, G., Valencia, V., Pullen, A., 2006, Detrital Zircon Geochronology by Laser Ablation Multicollector ICPMS at the Arizona LaserChron Center, in Olszewski, T., ed., *Geochronology: Emerging Opportunities: Paleontology Society Papers*, 12, 67-76.

Gehrels, G. E., Valencia, V., and Ruiz, J., 2008, Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation-multicollector-inductively coupled plasma-mass spectrometry: *Geochemistry, Geophysics, and Geosystems*, 9, 3.

Hann, H. P., 1987, Pegmatites from South Carpathians (*in Romanian*), Ed. Academiei, Bucureşti, 340 p.

Hann, H. P., 1995, Central South Carpathians: petrologic and structural investigations. *Romanian Journal of Petrology*, v. 76, 13-19.

Iancu, V., Seghedi, A., 2017, The South Carpathians: tectono-metamorphic units related to Variscan and Pan-African inheritance. *Geo-Eco-Marina* 23, 245-262.

Iancu, V., Maruntiu, M., Johan, V., Ledru, P., 1998. High-grade metamorphic rocks in the pre-Alpine nappe stack of the Getic-Supragetic basement (Median Dacides, South Carpathians, Romania). *Mineralogy and Petrology* 63, 173-198.

Iancu, V., Berza, T., Seghedi, A., Mărunțiu, M., 2005, Paleozoic rock assemblages incorporated in the South Carpathian Alpine thrust belt (Romania and Serbia): A review. *Geologica Belgica* 8, 48–68.

Krätner, H.G., 1997, Alpine and pre-Alpine terranes in the Romanian Carpathians and Apuseni Mountains, in: Papanikolaou, D. (Ed.), *Terrane Maps and Terrane Descriptions*. IGCP Project no. 276. *Annales Geologiques des Pays Helleniques*, Athens, pp. 331-400.

Liegeois, J.-P., Berza, T., Tatu, M., Duchesne, J.C., 1996, The Neoproterozoic Pan-African basement from the Alpine Lower Danubian nappe system (South Carpathians, Romania). *Precambrian Research* 80, 281-301.

Ledru, P., Coherie, A., Iancu, V., Mărunțiu, M., 1997, The gneissic units of the Median Dacides (Getic-Supragetic domain); an exotic segment of the European Variscides. *Romanian Journal of Mineralogy* 78, Supplement 1, 48.

Ludwig, K.R., 2008, *Isoplot/Ex 3.70. A Geochronological Toolkit for Microsoft Excel*. Berkeley Geochronological Center, Berkeley, Special publication No. 4., 76 pp.

Mațenco, L., 2017, Tectonics and Exhumation of Romanian Carpathians: Inferences from Kinematic and Thermochronological Studies. In: Radoane M., Vespremeanu-Stroe A. (eds) *Landform Dynamics and Evolution in Romania*. Springer Geography. Springer, Cham. [https://doi.org/10.1007/978-3-319-32589-7\\_2](https://doi.org/10.1007/978-3-319-32589-7_2).

Medaris, G., Ducea, M., Ghent, E., Iancu, V., 2003, Conditions and timing of high-pressure Variscan metamorphism in the South Carpathians, Romania. *Lithos* 70, 141–161.

Ménard, G., Molnar, P., 1988, Collapse of a Hercynian Tibetan Plateau into a late Palaeozoic Basin and Range Province in southern Europe, *Nature*, 334, 235-237.

Manecan, T., 1985, Studiul mineralogic si petrografic al pegmatitelor din bazinul Vaii Lotrului. PhD thesis, Univ. Bucuresti.

Pană, D., Erdmer, P., 1994, Alpine crustal shear zones and pre-Alpine basement terranes in the Romanian Carpathians and Apuseni Mountains, *Geology*, 22, 9, 807-810.

Poller, U., Uher, P., Broska, I., Plasienska, D., Janák, M., 2002, First Permian–Early Triassic zircon ages for tin-bearing granites from the Gemeric unit (Western Carpathians, Slovakia): connection to the post-collisional extension of the Variscan orogen and S-type granite magmatism *Terra Nova*, 14, 41-48.

Pullen, A., Ibáñez-Mejia, M., Gehrels, G. E., Giesler, D., and Pecha, M., 2018, Optimization of a Laser Ablation-Single Collector-Inductively Coupled Plasma-Mass Spectrometer (Thermo Element 2) for Accurate, Precise, and Efficient Zircon U-Th-Pb Geochronology: *Geochemistry, Geophysics, Geosystems*, 19, 10, 3689-3705, doi: 10.1029/2018gc007889.

Savu, H., Schuster, A.C., 1975, Harta geologică a R.S. România, sc. 1:50.000, foaia 107d, Voineasa, Inst.Geol.Geofiz., București.

Savu, H., Pavelescu, M., Stancu, I., Lupu, D., 1968, Harta geologică a R.S. România, sc. 1:200.000, foaia 26, Orastie, Inst. Geol. Geofiz., București

Săbău G., 1994, Lithostratigraphic and metamorphic correlations: A tentative way of exploring the early history of the Getic cristalline. *Rom. J. Petrology* 76, 19-128.

Săbău, G., Massone, H.J., 2003, Relationships among eclogite bodies and host rocks in the Lotru metamorphic suite (South Carpathians, Romania): Petrological evidence for multistage tectonic emplacement of eclogites in a medium-pressure terrain. *Int. Geol. Review* 45, 225-262.

Săndulescu, M., 1984, Geotectonics of Romania (*in Romanian*), București, Editura Tehnică, 336 pp.

Schmid, S., Bernoulli, D., Fügenschuh, B., Mațenco, L., Schefer, S., Schuster, R., Tischler, M., Ustaszewski, K., 2008, The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units, *Swiss Journal of Geosciences*, 101, 139-183, doi:10.1007/s00015-008-1247-3.

Schmid, S.M., Fügenschuh, B., Kounov, A., Mațenco, L., Nievergelt, P., Oberhänsli, R., Pleuger, J., Schefer, S., Schuster, R., Tomljenović, B., Ustaszewski, K., Van Hinsbergen, D.J.J., 2020, Tectonic units of the Alpine collision zone between Eastern Alps and western Turkey. *Gondwana Research* 78, 308–374.. doi:10.1016/j.gr.2019.07.005.

Stelea, I., 1994, Gneiss domes in the Sebeș-Lotru Series, Sebeș Mountains. *Rom. J. Petrology* 76, 143-148.

Stelea, I., 1999. Blastomilonitic Formations in Sebes Massif (*in Romanian*) PhD thesis, Fac. de Geol. si Geofiz. Bucuresti, Bucuresti, p. 132.

Stern, R.J., 1994, Arc assembly and continental collision in the Neoproterozoic East African orogen: Implications for the consolidation of Gondwanaland: *Annual Review of Earth and Planetary Sciences*, v. 22, p. 319–351. doi:10.1146/annurev.ea.22.050194.001535.

Stoica, A.M., Ducea, M.N., Roban, R.D., Jianu, D., 2016, Origin and evolution of the South Carpathians basement (Romania): a zircon and monazite geochronologic study of its Alpine sedimentary cover, *International Geology Review*, 58(4), 510-524, doi:10. 1080/00206814. 2015.1092097.

Uher, P., Broska, I., 1996, Post-orogenic Permian granitic rocks in the West Carpathian-Pannonian area: geochemistry, mineralogy and evolution. *Geologica Carpathica* 47, 311—321.

van Hinsbergen, D.J.J., Torsvik, T.H., Schmid, S.M., Mațenco, L.C., Maffione, M., Vissers, R.L.M., Gürer, D., Spakman, W., 2020, Orogenic architecture of the Mediterranean region and

kinematic reconstruction of its tectonic evolution since the Triassic. *Gondwana Research* 81, 79–229.. doi:10.1016/j.gr.2019.07.009.

von Raumer, J.F., Stampfli, G.M., 2008, The birth of the Rheic Ocean – Early Palaeozoic subsidence patterns and subsequent tectonic plate scenarios. *Tectonophysics* 461, 9–20, doi: 10.1016/j.tecto.2008.04.012.

von Raumer, J.F., Bussy, F., Schaltegger, U., Schulz, B., Stampfli, G.M., 2013, Pre-Mesozoic Alpine basements – Their place in the European Paleozoic framework: *Geological Society of America Bulletin*, v. 125, p. 89–108. doi:10.1130/B30654.1.









